

PRODUCTIVITY WITH PHYSICIAN EXTENDERS

tate not only the productivity of the practice, but the extent to which the physician must devote time or desires to be a manager or consultant for care rather than a direct provider of care.

REFERENCES

1. Yankauer A, Connelly JP, Feldman JJ: Physician productivity in the delivery of ambulatory care: Some findings from a survey of pediatricians. *Medical Care* 8:35-46, 1970
2. Bergman AB, Dassel SW, Wedgwood RJ: Time-motion study of practicing pediatricians. *Pediatrics* 38:254-263, 1966
3. Riddick FA, Bryan JB, Gershenson MI, et al: Use of allied health professionals in internists' offices. *Arch Intern Med* 127:924-931, 1971
4. Steiger WA, Yates LK: Patients; Needs and physician role. *Arch Environ Health* 18:806-810, 1969
5. Smith RA, Anderson JR, Akimotor JT: Increasing physician productivity and the hospitalization characteristics of practices using Medex—A progress report. *NW Med* 70:701-706, 1971
6. Duke Physician's Associate Program, Informational Pamphlet Series: Evaluations. Duke University School of Medicine, Department of Community Health Services, Durham, NC (undated)
7. Carlson CL, Athelstan GT: The physician's assistant, versions and diversions of a promising concept. *JAMA* 214:1855-1861, 1970
8. Uyeno DH: Health manpower systems: An application of simulation to the design of primary health care teams: *Management Science* 20:981-989, 1974
9. Golladay FL, Miller M, Smith KR: Allied health manpower strategies: Estimates of the potential gains from efficient task delegation. *Medical Care* 11:457-469, 1973
10. Lewis CE, Resnik BA: Nurse clinics and progressive ambulatory patient care. *N Engl J Med* 277:1236-1241, 1967
11. Sherman H, Reiffen B, Komraroff AL: Aids to the delivery of ambulatory medical care—IEEE Transactions on Biomedical Engr. *BME-20*:165-174, 1973
12. Sox HC, Sox CH, Tompkins RK: The training of physician's assistants, the use of a clinical algorithm system for patient care—Audit of performance and education. *N Engl J Med* 288:818-824, 1973
13. Merenstein JH, Wolfe H, Barker KM: The use of nurse practitioners in a general practice. *Med Care* 7:445-452, 1974
14. Couleman JL, Sheedy S: The role, training, and one-year's experience of a medical nurse practitioner. *Health Services Reports* 88:827-833, 1973

Managing Cardiac Tamponade

At Charity Hospital in New Orleans we have a trauma service that I think represents about 60 percent of all our surgical load. It's not unusual to see 10 or 11 instances of tamponade within a given month. When one sees a patient undergoing tamponade, don't try to make the diagnosis on a chest film. Almost invariably, in acute tamponade, you don't get the bottle-sized heart—that's chronic pericardial effusion. These patients often have a small heart. It may be an emergency film and it may not be adequate. You don't always see an elevated central venous pressure, particularly if the tamponade is severe enough so that cardiac output is impinged upon. In fact, the central venous pressure may even be zero at times. A high index of suspicion, really, is about the only thing you have to go on in a penetrating injury (gunshot wound or stab wound) usually in the region of the heart. If there's any question at all, don't hesitate to do a pericardiocentesis.

—THEODORE DRAPANAS, MD, *New Orleans*
Extracted from *Audio-Digest Surgery*, Vol. 22, No. 12, in the Audio-Digest Foundation's subscription series of tape-recorded programs. For subscription information: 1930 Wilshire Boulevard, Suite 700, Los Angeles, CA 90057.